

Table 4 Non-core vaccines for pet cats			
		Administration instructions	Clinically relevant comments for administration
Bordetella	INTRANASAL	For frequency and interval, follow label instructions	<ul style="list-style-type: none"> <li>❖ Not routinely used in pet cats</li> <li>❖ Provides incomplete protection</li> <li>❖ Use as part of a control program in a multi-cat household where infection is confirmed</li> <li>❖ Never administer parenterally</li> </ul>
	Attenuated live		
Chlamydia	PARENTERAL	For frequency and interval, follow label instructions	<ul style="list-style-type: none"> <li>❖ Provides incomplete protection</li> <li>❖ Use as part of a control program in a multi-cat household where infection is confirmed</li> <li>❖ Vaccination may be associated with a higher risk of adverse effects (lethargy, limb soreness, anorexia)<sup>32</sup></li> </ul>
	Inactivated		
	PARENTERAL	For frequency and interval, follow label instructions	<ul style="list-style-type: none"> <li>❖ Provides incomplete protection</li> <li>❖ Use as part of a control program in a multi-cat household where infection is confirmed</li> <li>❖ Vaccination may be associated with a higher risk of adverse effects (lethargy, limb soreness, anorexia)<sup>32</sup></li> </ul>
	Attenuated live		

Table 4 Non-core vaccines for pet cats